

AD-A103 254

REVERSIBLE HOMOGENEOUS CATALYSIS OF CARBON DIOXIDE  
HYDROGENATION/REDUCTIO. (U) CLARKSON UNIV POTSDAM NY  
DEPT OF CHEMISTRY S SCHREINER ET AL. 22 JUL 87 TR-1

1/1


UNCLASSIFIED

N00014-84-K-0658

F/G 7/2

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A183 254

OFFICE OF NAVAL RESEARCH

DTIC FILE COPY

12

Contract N00014-84-K-0658

R&T Code 413a003--01

Technical Report No. 1

Reversible Homogeneous Catalysis of Carbon Dioxide Hydrogenation/Reduction  
at Room Temperature and Low Pressures Mediated by a Platinum(0)-Diphosphine  
Cluster Complex in Solution



by

Serge Schreiner, James Y. Yu, and L. Vaska\*

Prepared for Publication

in the

Journal of the American Chemical Society

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Clarkson University  
Department of Chemistry  
Potsdam, NY 13676

July 22, 1987

DTIC  
ELECTE  
S AUG 13 1987 D  
E

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited.

87 8 103

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release and sale; its distribution is unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) No. 1			7a. NAME OF MONITORING ORGANIZATION Office of Naval Research		
6a. NAME OF PERFORMING ORGANIZATION Clarkson University		6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217		
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry (L. Vaska) Clarkson University Potsdam, NY 13676		8b. OFFICE SYMBOL (if applicable) 413	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-84-K-0658		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION ONR		10. SOURCE OF FUNDING NUMBERS			
8c. ADDRESS (City, State, and ZIP Code) See 7b		PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO
11. TITLE (Include Security Classification) Reversible Homogeneous Catalysis of Carbon Dioxide Hydrogenation/Reduction at Room Temperature and Low Pressures Mediated by a Platinum(0)-Diphosphine Cluster Complex in Solution					
12. PERSONAL AUTHOR(S) S. Schreiner, James Y. Yu, L. Vaska					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 1987/7/22	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION Submitted for publication in the Journal of the American Chemical Society					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Carbon Dioxide, Hydrogenation, Homogeneous Catalysis, N,N-dimethylformamide, Platinum, Phosphine Complex		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A platinum complex, $[Pt_2(\mu-PCP)_3]$ ( $PCP = (C_6H_5)_2PCH_2P(C_6H_5)_2$ ), catalyzes the synthesis of N,N-dimethylformamide (DMF) from $CO_2$ , $H_2$ and $(CH_3)_2NH$ in toluene solution at 25-100°C and total pressures of 0.96-114 atm of the three reactant gases. The turnover numbers, DMF/complex/day, range from 9 at 25°C and 0.96 atm to 1375 at 100°C and 114 atm. The homogeneous synthesis of DMF is readily reversible by either increasing the temperature, lowering the pressure of the reactants, or both, during the catalytic reaction. The formation of DMF appears to be dependent on the partial pressures of the individual reactants. Separate reactions of the three reactants with $[Pt_2(\mu-PCP)_3]$ at ambient conditions (25°C, 700 torr) give adducts with $(CH_3)_2NH$ , but not with $CO_2$ or $H_2$ . <i>(Keynote)</i>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTC USES			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL L. Vaska			22b. TELEPHONE (include Area Code) 315-268-2393/2389		22c. OFFICE SYMBOL

Reversible Homogeneous Catalysis of Carbon Dioxide Hydrogenation/Reduction at  
Room Temperature and Low Pressures Mediated by a Platinum(0)-Diphosphine Cluster  
Complex in Solution


Serge Schreiner, James Y. Yu, and L. Vaska\*

*Department of Chemistry, Clarkson University*


*Potsdam, New York 13676*

*Received*

Thermal reduction of carbon dioxide by dihydrogen catalyzed by transition metal complexes or supported metals has routinely been effected under forced conditions.<sup>1</sup> In homogeneous catalysis in solution, the reactions between CO<sub>2</sub>, H<sub>2</sub> and alcohols (ROH) or amines (R<sub>2</sub>NH) lead to formates<sup>2,3</sup> or formamides<sup>2,4</sup> (eq 1), and virtually all these conversions have been carried out at elevated




temperatures and pressures.<sup>2-4</sup> In heterogeneous catalysis by supported metals, the hydrogenation usually proceeds to methane<sup>5</sup> or methanol<sup>6</sup> (eq 2), and the



processes invariably require high temperatures.<sup>5,6</sup> A notable exception to forced conditions is the reduction of bicarbonate to formate in aqueous solution by supported palladium catalysts at ambient conditions.<sup>7</sup>

We wish to report that a diplatinum complex, [Pt<sub>2</sub>(μ-PCP)<sub>3</sub>] (PCP = (C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>PCH<sub>2</sub>P(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>),<sup>8</sup> is an effective catalyst precursor for reaction (3) at





mild conditions. The catalysis takes place in homogeneous toluene solution at room temperature and is clearly observable even under a total pressure of less than one atm of the three reactant gases (eq 3). Furthermore, the reaction is readily reversible, an unusual observation in homogeneous catalysis by metal complexes.

Some pertinent results are summarized in Table I. Two sets of experiments were carried out: (a) under total pressures of ca. 1 atm (entries 1,2), and (b) at elevated and varying pressures (entries 3,4). At 25°C and ca. 1 atm, the equilibrium (eq 3) was attained in about 24 h. When the same solution was subsequently warmed to 50°C, the yield of DMF *decreased* substantially (entry 1b), indicating that the reverse reaction (eq 3) had taken place. A separate experiment, commencing at 70°C, gave no DMF or any other product for 5 days (entry 2).

Further evidence for the reversibility of the catalysis is obtained from reactions at higher pressures. Entries 3 and 4 in Table I refer to two runs with pressure-temperature variations in the sequences shown (a+b). Experiment 3a,b demonstrates that the catalytic synthesis of DMF is reversed (eq 3) by simply lowering the initially applied total pressure<sup>9</sup> at room temperature. And the combined effects of temperature and pressure<sup>9</sup> on the catalytic reversibility of the system are shown by entry 4a,b,c.

Although thermochemical values for reaction (3) in toluene are not available, the data in other environments (eq 4,5) indicate that there is a





$$\Delta H^\circ = -36.3, \Delta G^\circ = -0.75 \quad (5)^{10}$$

delicate balance between the reactants and products which explains the observed ready reversibility thermodynamically. Kinetically, the facile catalysis with an apparently low activation energy seems to derive from the structural<sup>8d</sup> features of the catalyst precursor,  $[\text{Pt}_2(\mu\text{-PCP})_3]$ . The complex displays two terminal vacant coordination sites and a third one for insertion into the Pt-Pt bond. With CO, a fast reversible reaction gives a dicarbonyl complex,  $[(\text{CO})\text{Pt}(\mu\text{-PCP})_3\text{Pt}(\text{CO})]$  ( $\nu_{\text{CO}} = 1940 \text{ cm}^{-1}$ ), and with oxygen, a peroxo species is produced with this possible formulation,  $[\text{Pt}(\eta^2\text{-}\mu\text{-O}_2)(\mu\text{-PCP})_3\text{Pt}]$  ( $\nu_{\text{O}_2} = 827 \text{ cm}^{-1}$ ).<sup>8b</sup> Measurements of gas uptake of the three reactants (eq 3) individually by  $[\text{Pt}_2(\mu\text{-PCP})_3]$  in toluene solutions at 25°C and 700 torr showed that dimethylamine forms adducts of indefinite compositions, but there is, perhaps surprisingly, no detectable reaction with carbon dioxide or hydrogen *under these conditions*.<sup>8b</sup> Yet, the activation of both  $\text{CO}_2$  and  $\text{H}_2$  by  $[\text{Pt}_2(\mu\text{-PCP})_3]$  under low pressures is implicated by the catalytic results (entry 1, Table I).<sup>11</sup>

The selectivity of the DMF synthesis was found to be about 90-95%. The only other catalytic product was trimethylamine. It should also be noted that in all experiments, the final reaction mixture contained dimethylammonium carbamate,  $[(\text{CH}_3)_2\text{NH}_2]^+[(\text{CH}_3)_2\text{NC}(\text{O})\text{O}]^-$ , which is spontaneously (non-catalytically) formed from  $\text{CO}_2$  and  $(\text{CH}_3)_2\text{NH}$ .<sup>13</sup> Considering all these complications, including the different individual reactivities of the three catalytic reactants toward  $[\text{Pt}_2(\mu\text{-PCP})_3]$ ,<sup>9</sup> it is premature to speculate about the mechanism of the overall catalysis (eq 3).<sup>11</sup>

Acknowledgment. This work was supported in part by the Office of Naval Research. We thank P.B. Kaufman and J.K. Hartikainen for helpful experimental assistance.

Table I. Reversible Catalytic Formation of N,N-dimethylformamide (DMF) from CO<sub>2</sub>, H<sub>2</sub> and (CH<sub>3</sub>)<sub>2</sub>NH in Toluene (50 mL) Solution Mediated by [Pt<sub>2</sub>(μ-PCP)<sub>3</sub>] (3-9 x 10<sup>-5</sup> mol) (eq 3)<sup>a</sup>

entry <sup>b</sup>	reaction temp, °C	total pressure, atm <sup>c</sup>	turnover, <sup>d</sup> DMF/Pt <sub>2</sub> /day
1a	25	0.96	8.7
1b	50	0.83	-5.7
2	70	0.75	0
3a	25	101	9.3
3b	25	5.4	-8.2
4a	100	114	1375
4b	100	1.0	-409
4c	25	92	486

<sup>a</sup>Experiments were carried out for periods lasting from one to several days and by using a gas burette-manometer system (entries 1,2) or a Parr pressure vessel (entries 3,4). Each entry represents a 24-h reaction period. Samples were periodically withdrawn from the reactors and analyzed by IR and GC methods. DMF was quantitatively determined on a pre-calibrated Carbowax 20 M + KOH GC column. Blank runs under the same conditions, i.e., with all the reactants present except the catalyst precursor (eq 3), gave no products. <sup>b</sup>Each number refers to a separate experiment, the letters (a,b,c) refer to sequential variation of conditions within that experiment. <sup>c</sup>At reaction temperature. The initial pressures of the reactants applied at 25°C ranged as follows: entries 1,2: CO<sub>2</sub>, 0.20-0.27; H<sub>2</sub>, 0.33-0.51; (CH<sub>3</sub>)<sub>2</sub>NH, 0.14-0.17; entries 3,4: CO<sub>2</sub>, 13.9-87.4; H<sub>2</sub>, 20.4-79.6; (CH<sub>3</sub>)<sub>2</sub>NH, 1.02 atm. <sup>d</sup>The yield of DMF (mol)/[Pt<sub>2</sub>(μ-PCP)<sub>3</sub>] (mol, introduced initially)/day (reaction period). The



Footnote d to Table I contd.

negative turnover numbers indicate the rates of the reverse reaction (eq 3),  
i.e., the disappearance of DMF produced during the preceding reaction period.

## References and Notes

(1) Thermal catalysis of CO<sub>2</sub> reduction is to be distinguished from the systems which operate on supplied photochemical or electrochemical energy and usually at ambient temperatures and pressures. See, for example, these recent papers and the references quoted therein: (a) Kutal, C; Corbin, A.J.; Ferraudi, G. *Organometallics* 1987, 6, 553-557. (b) Ishida, H.; Tanaka, K.; Tanaka, T. *Organometallics* 1987, 6, 181-186. (c) DuBois, D.L.; Miedaner, A. *J. Am. Chem. Soc.* 1987, 109, 113-117.

(2) For recent reviews on CO<sub>2</sub> activation by metal complexes and related reactions, see: (a) Inoue, S.; Yamasaki, N., Eds. *Organic and Bio-organic Chemistry of Carbon Dioxide*; John Wiley and Sons: New York, 1982. (b) Behr, A. In *Catalysis of C<sub>1</sub> Chemistry*; Keim, W., Ed.; D. Reidel: Dordrecht, Holland, 1983; pp. 169-217. (c) Darensbourg, D.J.; Kudarowski, R.A. *Adv. Organometal. Chem.* 1983, 22, 129-168.

(3) Darensbourg, D.J.; Ovalles, C. *J. Am. Chem. Soc.* 1987, 109, 3330-3336, and references therein.

(4) (a) Haynes, P.; Slauch, L.H.; Kohnle, J.F. *Tetrahedron Lett.* 1970, 365-368. (b) Kudo, K.; Phala, H.; Sugita, N.; Takezaki, Y. *Chem. Lett.* 1977, 1495-1496. (c) Phala, H.; Kudo, K.; Sugita, N. *Bull. Inst. Chem. Res., Kyoto Univ.* 1981, 59, 88-105.

(5) (a) Erdöhelyi, A.; Pásztor, M.; Solymosi, F. *J. Catalysis* 1986, 98, 166-177. (b) Darensbourg, D.J.; Ovalles, C. *Inorg. Chem.* 1986, 25, 1603-1609; and references cited in these two papers.

(6) (a) Denise, B.; Sneed, R.P.A. *J. Mol. Catalysis* 1982, 17, 359-366. (b) Iizuka, T.; Kojima, M.; Tanabe, K. *J. Chem. Soc., Chem. Commun.* 1983, 638-639. (c) He, M.-Y.; White, J.M. *J. Mol. Catalysis* 1985, 30, 415-430. (d) Ref. 5a; and references quoted in the four papers cited above.

(7) Stalder, J.C.; Chao, S.; Summers, D.P.; Wrighton, M.S. *J. Am. Chem.*

*Soc.* 1983, 105, 6318-6320.

(8) (a) Chin, C.-S.; Sennett, M.S.; Wier, P.J.; Vaska, L. *Inorg. Chim. Acta* 1978, 31, L443-L444. (b) Chin, C.-S.; Wier, P.J.; Sennett, M.S.; Kim, S.-H.; Vaska, L., unpublished results. (c) Grossel, M.C.; Brown, M.P.; Nelson, C.D.; Yavari, A.; Kallas, E.; Moulding, R.P.; Seddon, K.R. *J. Organometal. Chem.* 1982, 232, C13-C16. (d) Manojlović-Muir, L.; Muir, K.W.; Grossel, M.C.; Brown, M.P.; Nelson, C.D.; Yavari, A.; Kallas, E.; Moulding, R.P.; Seddon, K.R. *J. Chem. Soc., Dalton Trans.* 1986, 1955-1963.

(9) It is actually the applied partial pressures of CO<sub>2</sub>, H<sub>2</sub> and (CH<sub>3</sub>)<sub>2</sub>NH, and the consequent solubilities (concentrations) of the gases, which appear to determine the yield of DMF. The individual pressures were varied from experiment to experiment (footnote c to Table I). Thus, runs at similar total pressures (and temperatures) are not necessarily comparable. For example, it has been found that a relatively high CO<sub>2</sub> pressure lowers the yield due, probably, to suppressing the concentration of the active catalyst (cf. entries 3a and 4c, with P<sub>CO<sub>2</sub></sub> = 87.4 and 13.9 atm, respectively).

(10) Calculated from the ΔH<sub>f</sub><sup>0</sup> and ΔG<sub>f</sub><sup>0</sup> values of the reactants and products: (a) *J. Chem. Phys. Ref. Data* 1982, 11, Supplement No. 2. (b) Dean, J.A., Ed. *Lange's Handbook of Chemistry*, 13th Ed.; McGraw Hill: New York, 1985. (c) For notation of standard states, see ref. 12a.

(11) It is hoped that *in situ* catalytic studies under elevated pressures by using cylindrical internal reflectance coupled with an FTIR spectrometer<sup>12</sup> will shed some light on these and other mechanistic questions.

(12) Moser, W.R.; Chossen, J.E.; Wang, A.W.; Krouse, S.A. *J. Catalysis* 1985, 95, 21-32.

(13) A common reaction between CO<sub>2</sub> and amines or ammonia; see, for example: Wright, H.B.; Moore, M.B. *J. Am. Chem. Soc.* 1948, 70, 3865-3866, and references quoted.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

END

9-87

Dtic